

(51) International Patent Classification 5 : C07K 7/06, 7/08, 7/10 A61K 37/02		A1	(11) International Publication Number: WO 93/01211 (43) International Publication Date: 21 January 1993 (21.01.93)
(21) International Application Number: PCT/AU92/00332 (22) International Filing Date: 3 July 1992 (03.07.92) (30) Priority data: PK 7097                      5 July 1991 (05.07.91)                      AU PK 7924                      22 August 1991 (22.08.91)                      AU (71) Applicant (for all designated States except US): PEPTIDE TECHNOLOGY LIMITED [AU/AU]; 4-10 Inman Road, Dee Why, NSW 2099 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only) : RATHJEN, Deborah, Ann [AU/AU]; 4 Eddy Street, Thornleigh, NSW 2120 (AU). WIDMER, Fred [CH/AU]; 35 Anzac Avenue, Ryde, NSW 2112 (AU). GRIGG, Geoffrey, Walter [AU/AU]; 352 Burns Bay Road, Lane Cove, NSW 2066 (AU). MACK, Philip, On-Lok [CN/AU]; 22 Ethel Street, Hornsby, NSW 2077 (AU).		(74) Agent: F.B. RICE & CO; 28A Montague Street, Balmain, NSW 2041 (AU). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published With international search report.	
(54) Title: PEPTIDE WHICH ABROGATES TNF AND/OR LPS TOXICITY			
<p>VRSSSRTPSD<sub>10</sub>KPVAHVVANP<sub>20</sub>QAEGQLQWLN<sub>30</sub>RRA</p> <p>NALLANG<sub>40</sub>VELRDNQLVW<sub>50</sub>PSEGLYLIYS<sub>60</sub>QVLFKGQGCP<sub>70</sub>STHVLL</p> <p>THTI<sub>80</sub>SRIAVSYQTK<sub>90</sub>VNLLSAIKSP<sub>100</sub>CQRETREGAE<sub>110</sub>AKPWYEPI</p> <p>YL<sub>120</sub>GGVFQLEKGD<sub>130</sub>RLSAEINRPD<sub>140</sub>YLDFAESGQV<sub>150</sub>YFGIIAL<sub>157</sub></p>			
(57) Abstract			
<p>The present invention provides peptides which have the ability to abrogate TNF toxicity and/or LPS toxicity. The present invention further relates to compositions including these peptides as the active ingredient and methods of anti-inflammatory treatment involving the administration of this composition. The peptides of the present invention are based primarily on residue 1 to 26 of human TNF.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NI	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland			SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		
ES	Spain	MG	Madagascar		

- 1 -

PEPTIDE WHICH ABROGATES TNF AND/OR LPS TOXICITYField of the Invention

The present invention relates to a group of peptides which have the ability to abrogate TNF toxicity and/or LPS toxicity. The present invention further relates to compositions including this peptide as the active ingredient and methods of anti-inflammatory treatment involving the administration of this composition.

Background of the Invention

Many of the clinical features of septicemic shock induced by Gram-negative bacteria which have lipopolysaccharide (LPS) in their cell walls may be reproduced in animals by the administration of LPS. This induces prompt severe metabolic and physiological changes which can lead to death. Associated with the injection of LPS is the extensive production of tumour necrosis factor alpha (TNF). Many of the effects of LPS injection or indeed of Gram-negative bacteria can be reproduced by TNF. Thus, mice injected with recombinant human TNF develop piloerection of the hair (ruffling), diarrhoea, a withdrawn, unkempt appearance and die if sufficient amounts are given. Rats treated with TNF become hypotensive, tachypneic and die of sudden respiratory arrest (Tracey et al., 1986 Science 234, 470). Severe acidosis, marked haemoconcentration and biphasic changes in blood glucose concentration were also observed. Histopathology revealed severe leukostasis in the lungs, haemorrhagic necrosis in the adrenals, pancreas and other organs and tubular necrosis of the kidneys. All these changes were prevented if the animals were pretreated with a neutralizing monoclonal antibody against TNF.

The massive accumulation of neutrophils in the lungs of TNF-treated animals reflects the activation of neutrophils by TNF. TNF causes neutrophil degranulation, respiratory burst, enhanced antimicrobicidal and

- 2 -

anti-tumour activity (Klebanoff et al., 1986 J. Immunol. 136, 4220; Tsujimoto et al., 1986 Biochem Biophys Res Commun 137, 1094). Endothelial cells are also an important target for the expression of TNF toxicity. TNF diminishes the anticoagulant potential of the endothelium, inducing procoagulant activity and down regulation of the expression of thrombomodulin (Stern and Nawroth, 1986 J Exp Med 163, 740).

10 TNF, a product of activated macrophages produced in response to infection and malignancy, was first identified as a serum factor in LPS treated mice which caused the haemorrhagic necrosis of transplantable tumours in murine models and was cytotoxic for tumour cells in culture (Carswell et al., 1975 PNAS 72, 3666; Nelson et al., 1975 Nature 258, 731). Cachexia is a common symptom of advanced malignancy and severe infection. It is characterised by abnormal lipid metabolism with hypertriglyceridemia, abnormal protein and glucose metabolism and body wasting. Chronic administration of TNF (also known as cachectin in the early literature) to mice causes anorexia, weight loss and depletion of body lipid and protein within 7 to 10 days (Cerami et al., 1985 Immunol Lett 11, 173; Fong et al., 1989 J Exp Med 170, 1627). These effects were reduced by concurrent administration of antibodies against TNF. Although TNF has been measured in the serum of patients with cancer and chronic disease associated with cachexia the results are inconclusive since large differences in TNF levels have been reported. These may be due to the short half-life of TNF (6 minutes), differences in TNF serum binding protein, or true differences in TNF levels in chronic disease states.

30 TNF $\alpha$ , as a mediator of inflammation, has been implicated in the pathology of other diseases apart from toxic shock and cancer-related cachexia. TNF has been

35

- 3 -

measured in synovial fluid in patients with both  
rheumatoid and r active arthritis and in the serum of  
patients with rheumatoid arthritis (Saxne t al., 1988  
Arthrit. Rheumat. 31, 1041). Raised levels of TNF have  
5 been detected in renal transplant patients during acute  
rejection episodes (Maury and Teppo 1987 J. Exp Med 166,  
1132). In animals TNF has been shown to be involved in  
the pathogenesis of graft versus host disease in skin and  
gut following allogeneic marrow transplantation.

10 Administration of a rabbit anti-murine TNF was  
demonstrated to prevent the histological changes  
associated with graft versus host disease and reduced  
mortality (Piquet et al., 1987 J Exp Med 166, 1280).

TNF has also been shown to contribute significantly  
15 to the pathology of malaria (Clark et al., 1987; Am. J.  
Pathol. 129: 192-199). Further, elevated serum levels of  
TNF have been reported in malaria patients (Scuderi  
et al., 1986; Lancet 2: 1364-1365). TNF may also  
contribute to the brain pathology and consequent dementia  
20 observed in late stage HIV infections (Grimaldi et al Ann  
Nevrol 29 : 21)

The peptides encompassed in the present invention do  
not necessarily interfere directly with the bio-synthetic  
mechanisms of the disease-causing component. As will be  
25 described below in the experimental data the mechanism  
behind the alleviating effect of the peptides is to be  
found in the modulation of the different cytokines  
produced by activated cells belonging to the cell-lines  
encompassing the immune defence. This modulation of  
30 cytokines is not limited to TNF but may also be valid for  
the whole range of interleukins, from interleukin-1 to  
interleukin 10. LPS, a known component of bacteria  
important in inducing major inflammatory response was used  
as a model. LPS binds to receptors on neutrophils,  
35 monocytes, endothelial cells and machrophages, which

- 4 -

consequently become activated and start production of IL-1 and TNF and other cytokin s, thus starting the inflammatory cascade. One parameter used to measure the effect of LPS is the concentration of blood glucose, which will normally decrease on exposure to TNF or LPS.

LPS normally combines with LPS-Binding-Protein (LBP) and exerts its dramatic effect through the CD14 receptor. The activation of the CD14 molecule by LPS results in TNF production by leucocytes. It is believed that the peptides of the present invention which abrogate LPS toxicity may exert their effect by interacting with the CD14 molecule and thus inhibit LPS binding.

The peptides identified by the present inventors which have the ability to abrogate TNF and/or LPS toxicity resemble peptide sequences found in the amino terminal of TNF $\alpha$ . Other investigators have also considered this area of the TNF $\alpha$  molecule but with little success in obtaining biologically active peptides.

In this regard attention is drawn to Canadian patent application Nos 2005052 and 2005056 in the name of BASF AG. Both these applications claim a wide range of peptide sequences and, by selecting appropriate alternatives it can be seen that application No 2005052 is directed toward the peptide sequence 7-42 of TNF $\alpha$  whilst application No 2005056 is directed toward amino acid sequence 1 to 24 of TNF $\alpha$ . Whilst each of these applications claim a broad range of peptide sequences it is noted that there is no indication as to what, if any, biological activity the claimed peptides may possess. Indeed there is no demonstration that any of the produced peptide have any biological activity. In contrast, the present inventors have produced a range of peptides which have specific activities in that they abrogate TNF and/or LPS toxicity.

#### Summary of the Invention

In a first aspect the present invention consists in a linear or cyclic peptide of the g n ral formula:-

- 5 -

$$X_1-X_2-X_3-X_4-X_5-X_6-X_7-X_8-X_9$$

in which

- $X_1$  is null, Cys or  $R_1$   
 $X_2$  is null, Cys,  $R_1$  or  $A_1-A_2-A_3-A_4-A_5$   
 5 in which  $A_1$  is Val or Ile or Leu or Met or His  
 $A_2$  is Arg or Cys or His  
 $A_3$  is Ser or Thr or Ala  
 $A_4$  is Ser or Thr or Ala  
 $A_5$  is Ser or Thr or Ala  
 10  $X_3$  is Cys,  $R_1$  or  $A_6-A_7$   
 in which  $A_6$  is Arg or Cys or His or Absent  
 $A_7$  is Thr or Ser or Ala  
 $X_4$  is Cys,  $R_1$  or  $A_8-A_9$   
 in which  $A_8$  is Pro or an  $\alpha$ -alkylamino acid  
 15  $A_9$  is Ser or Thr or Ala  
 $X_5$  is Cys,  $R_1$  or  $A_{10}$   
 in which  $A_{10}$  is Asp or Ala or Cys or Glu or Gly  
 or Arg or His  
 $X_6$  is Cys,  $R_2$  or  $A_{11}-A_{12}-A_{13}$   
 20 in which  $A_{11}$  is absent or Cys or Arg or His or  
 Asp or Glu  
 $A_{12}$  is Pro or an  $\alpha$ -alkylamino acid  
 $A_{13}$  is Val or Ile or Phe or Tyr or Trp  
 or His or Leu or His or Met  
 25  $X_7$  is null, Cys,  $R_2$  or  $A_{14}-A_{15}$   
 in which  $A_{14}$  is Ala or Val or Gly or Ile or Phe  
 or Trp or Tyr or Leu or His or Met  
 $A_{15}$  is absent or His or Arg or Glu or  
 Asa or Ala or Lys or Asp or Phe or Tyr or  
 30 Trp or Glu or Gln or Ser or Thr or Gly  
 $X_8$  is null, Cys,  $R_2$ ,  $A_{16}$ ,  $A_{16}-A_{17}$ ,  $A_{16}-A_{17}-A_{18}$  or  
 $A_{16}-A_{17}-A_{18}-A_{19}-A_{20}-A_{21}-A_{22}-A_{23}-A_{24}-A_{25}-A_{26}$   
 in which  $A_{16}$  is Val or Ile or Leu or Met or His  
 $A_{17}$  is Val or Ile or Leu or M t or His  
 35  $A_{18}$  is Ala or Gly

- 6 -

- A<sub>19</sub> is Asp or Glu  
A<sub>20</sub> is Pro or an N $\alpha$ -alkylamino acid  
A<sub>21</sub> is Gln or Asn  
A<sub>22</sub> is Ala or Gly  
5 A<sub>23</sub> is Glu or Asp  
A<sub>24</sub> is Gly or Ala  
A<sub>25</sub> is Gln or Asn  
A<sub>26</sub> is Leu or Ile or Val or Met or His  
X<sub>9</sub> is null, Cys or R<sub>2</sub>  
10 R<sub>1</sub> is R-CO, where R is H, straight, branched or  
cyclic alkyl up to C20, optionally containing double  
bonds and/or substituted with halogen, nitro, amino,  
hydroxy, sulfo, phospho or carboxyl groups (which may  
be substituted themselves), or aralkyl or aryl  
15 optionally substituted as listed for the alkyl and  
further including alkyl, or R<sub>1</sub> is glycosyl,  
nucleosyl, lipoyl or R<sub>1</sub> is an L- or D- $\alpha$  amino acid  
or an oligomer thereof consisting of up to 5 residues  
R<sub>1</sub> is absent when the amino acid adjacent  
20 is a desamino-derivative.  
R<sub>2</sub> is  
-NR<sub>12</sub>R<sub>13</sub>, wherein R<sub>12</sub> and R<sub>13</sub> are  
independently H, straight, branched or cyclic alkyl,  
aralkyl or aryl optionally substituted as defined for  
25 R<sub>1</sub> or N-glycosyl or N-lipoyl  
-OR<sub>14</sub>, where R<sub>14</sub> is H, straight, branched or  
cyclic alkyl, aralkyl or aryl, optionally substituted  
as defined for R<sub>1</sub>  
-O-glycosyl, -O-lipoyl or  
30 - an L- or D- $\alpha$ -amino acid or an oligomer thereof  
consisting of up to 5 residues  
or R<sub>2</sub> is absent, when the adjacent amino acid is a  
decarboxy derivative of cysteine or a homologue  
thereof or the peptide is in a N-C cyclic form.  
35 with the proviso that:



- 7 -

- when  $X_6$  is Cys or  $R_2$  then  $X_5$  is  $A_{10}$ ,  $X_4$  is  $A_8-A_9$ ,  
 $X_3$  is  $A_6-A_7$  and  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$   
 when  $X_5$  is Cys or  $R_1$  then  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  
 $A_{14}-A_{15}$ ,  $X_8$  is  $A_{16}-A_{17}-A_{18}$  and  $A_{11}$  is absent  
 5 when  $X_4$  is Cys or  $R_1$  then  $X_5$  is  $A_{10}$ ,  $X_6$  is  
 $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  $A_{14}-A_{15}$  and  $X_8$  is  
 $A_{16}-A_{17}-A_{18}$   
 when  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$  then  $X_8$  is not  $A_{16}$   
 when  $X_1$  is null,  $X_2$  is Cys or  $R_1$ ,  $X_3$  is  $A_6-A_7$ ,  $X_4$  is  
 10  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  
 $A_{14}-A_{15}$  and  $X_8$  is  $A_{16}$  then  $A_{16}$  is not D-His.  
 $X_1$  is always and only null when  $X_2$  is  $R_1$ , Lys or Null  
 $X_2$  is always and only null when  $X_3$  is Cys or  $R_1$   
 $X_3$  is always and only null when  $X_6$  is Cys or  $R_2$   
 15  $X_7$  is always and only null when  $X_7$  is Cys,  $R_2$  or Null  
 $X_8$  is always and only null when  $X_8$  is Cys,  $R_2$  or Null  
 $X_9$  is always and only null when  $X_8$  is Cys,  $R_2$  or Null  
 when  $X_1$  and  $R_2$  are null,  $X_3$  is  $R_1$ ,  $X_4$  is  
 20  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$   
 is  $A_{14}-A_{15}$ ,  $X_8$  is  $R_2$  and  $A_{14}$  is Ala and  $A_{15}$  is  
 absent then  $R_1$  is acetyl and  $R_2$  is  $NH_2$ .

The amino acids may be D or L isomers, however  
 generally the peptide will primarily consist of L-amino  
 25 acids.

In a second aspect the present invention consists in  
 a pharmaceutical composition for use in treating subjects  
 suffering from toxic effects of TNF and/or LPS, the  
 composition comprising a therapeutically effective amount  
 30 of a peptide of the first aspect of the present invention  
 and a pharmaceutically acceptable sterile carrier.

In a third aspect the present invention consists in a  
 method of treating a subject suffering from the toxic  
 effects of TNF and/or LPS, the method comprising  
 35 administering to the subject a therapeutically effective  
 amount of the composition of the second aspect of the  
 present invention.

- 8 -

In a preferred embodiment of the present invention

- ~ X<sub>1</sub> is H, X<sub>2</sub> is A<sub>1</sub>-A<sub>2</sub>-A<sub>3</sub>-A<sub>4</sub>-A<sub>5</sub>, X<sub>3</sub> is  
A<sub>6</sub>-A<sub>7</sub>, X<sub>4</sub> is A<sub>8</sub>-A<sub>9</sub>, X<sub>5</sub> is A<sub>10</sub>, X<sub>6</sub> is  
A<sub>11</sub>-A<sub>12</sub>-A<sub>13</sub>, X<sub>7</sub> is A<sub>14</sub>-A<sub>15</sub>, X<sub>8</sub> is  
5 A<sub>16</sub>-A<sub>17</sub>-A<sub>18</sub> and X<sub>9</sub> is OH.

In a further preferred embodiment of the present invention X<sub>1</sub> is null, X<sub>2</sub> is H or Ac, X<sub>3</sub> is

- A<sub>6</sub>-A<sub>7</sub>, X<sub>4</sub> is A<sub>8</sub>-A<sub>9</sub>, X<sub>5</sub> is A<sub>10</sub>, X<sub>6</sub> is  
A<sub>11</sub>-A<sub>12</sub>-A<sub>13</sub>, X<sub>7</sub> is A<sub>14</sub>-A<sub>15</sub>, X<sub>8</sub> is  
10 A<sub>16</sub>-A<sub>17</sub>-A<sub>18</sub> and X<sub>9</sub> is OH or NH<sub>2</sub>.

In a further preferred embodiment of the present invention X<sub>1</sub> is H, X<sub>2</sub> is A<sub>1</sub>-A<sub>2</sub>-A<sub>3</sub>-A<sub>4</sub>-A<sub>5</sub>,

- X<sub>3</sub> is A<sub>6</sub>-A<sub>7</sub>, X<sub>4</sub> is A<sub>8</sub>-A<sub>9</sub>, X<sub>5</sub> is A<sub>10</sub>, X<sub>6</sub>  
is OH and X<sub>6</sub>, X<sub>7</sub> and X<sub>8</sub> are null.  
15 In a further preferred embodiment of the present invention the peptide is selected from the group consisting of:-

- Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala  
-His-Val-Val-Ala;  
20 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
Arg-Thr-Pro-Ser-Ala-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
Arg-Thr-Pro-Ser-Lys-Asp-Pro-Val-Ala-His-Val-Val-Ala;  
Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala  
-Arg-Val-Val-Ala;  
25 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala  
-Gln-Val-Val-Ala;  
Ac-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-NH<sub>2</sub>;  
Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Ala-Val;  
Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Lys-Val;  
30 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val;  
Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val;  
Pro-Ser-Asp-Lys-Pro-Val-Ala-His;  
Pro-Ser-Asp-Lys-Pro-Val;  
Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-  
35 Val-His-Val-Val-Ala;

- 9 -

Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala-Asn  
 -Pro-Gln-Ala-Glu-Gly-Gln-Leu;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp;  
 Ac-Pro-Ser-Asp-Lys-Pro-Val-Ala-NH<sub>2</sub>;  
 5 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Asp-Val;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-  
 Ala-His-Val-Val-Ala-Asn-Pro-Gln-Ala-Glu-Gly-Gln-Leu;  
 Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val;  
 10 Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Pro-Sir-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Pro-Val-Ala-His-Val-Val-Ala; and  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Val-His-Val.

The composition and method of the present invention  
 15 would be expected to be useful as an anti-inflammatory  
 agent in a wide range of disease states including toxic  
 shock, adult respiratory distress syndrome,  
 hypersensitivity pneumonitis, systemic lupus  
 erythromatosis, cystic fibrosis, asthma, bronchitis, drug  
 20 withdrawal, schistosomiasis, sepsis, rheumatoid arthritis,  
 acquired immuno-deficiency syndrome, multiple sclerosis,  
 leperosy, malaria, systemic vasculitis, bacterial  
 meningitis, cachexia, dermatitis, psoriasis, diabetes,  
 neuropathy associated with infection or autoimmune  
 25 disease, ischemia/reperfusion injury, encephalitis,  
 Guillame Barre Syndrome, atherosclerosis, chronic fatigue  
 syndrome, TB, other viral and parasitic diseases, OKT3  
 therapy, and would be expected to be useful in conjunction  
 with radiation therapy, chemotherapy and transplantation,  
 30 to ameliorate the toxic effects of such treatments or  
 procedures.

As the peptide of the present invention suppresses  
 activation of neutrophils the composition and method of  
 the present invention may also be useful in the treatment  
 35 of diseases with an underlying element of local, systemic,  
 acute or chronic inflammation. In general, it is believed

- 10 -

the composition and method of the present invention will be useful in treatment of any systemic or local infection leading to inflammation.

The peptides of the present invention may also be administered in cancer therapy in conjunction with cytotoxic drugs which may potentiate the toxic effects of TNF $\alpha$  (Watanabe et al., 1988; Immunopharmacol. Immunotoxicol. 10: 117-127) such as vinblastin, acyclovir, interferon alpha, cyclosporin A, IL-2, actinomycin D, adriamycin, mitomycin C, AZT, cytosine arabinoside, daunorubicin, cis-platin, vincristine, 5-fluorouracil and bleomycin; in cancer patients undergoing radiation therapy; and in AIDS patients (or others suffering from viral infection such as viral meningitis, hepatitis, herpes, green monkey virus etc.) and in patients receiving immunostimulants such as thymopentin and muramyl peptides or cytokines such as IL-2 and GM-CSF. In this use peptides of the present invention will serve to abrogate the deleterious effects of TNF $\alpha$

It will be appreciated by those skilled in the art that a number of modifications may be made to the peptide of the present invention without deleteriously effecting the biological activity of the peptide. This may be achieved by various changes, such as insertions, deletions and substitutions (e.g., sulfation, phosphorylation, nitration, halogenation), either conservative or non-conservative (e.g., W-amino acids, desamino acids) in the peptide sequence where such changes do not substantially altering the overall biological activity of the peptide. By conservative substitutions the intended combinations are:-

G, A; V, I, L, M; D, E; N, Q; S, T; K, R, H; F, Y, W, H; and P, N $\alpha$ -alkylamino acids.

It may also be possible to add various groups to the peptide of the present invention to confer advantages such as increased potency or extended half-life in vivo,

- 11 -

without substantially altering the overall biological activity of the peptide.

The term peptide is to be understood to embrace peptide bond replacements and/or peptide mimetics, i.e. pseudopeptides, as recognised in the art (see for example: Proceedings of the 20th European Peptide Symposium, ed. G. Jung. E. Bayer, pp. 289-336, and references therein), as well as salts and pharmaceutical preparations and/or formulations which render the bioactive peptide(s) particularly suitable for oral, topical, nasal spray, ocular pulmonary, I.V., subcutaneous, as the case may be, delivery. Such salts, formulations, amino acid replacements and pseudopeptide structures may be necessary and desirable to enhance the stability, formulation, deliverability (e.g., slow release, prodrugs), or to improve the economy of production, and they are acceptable, provided they do not negatively affect the required biological activity of the peptide.

Apart from substitutions, three particular forms of peptide mimetic and/or analogue structures of particular relevance when designating bioactive peptides, which have to bind to a receptor while risking the degradation by proteinases and peptidases in the blood, tissues and elsewhere, may be mentioned specifically, illustrated by the following examples: Firstly, the inversion of backbone chiral centres leading to D-amino acid residue structures may, particularly at the N-terminus, lead to enhanced stability for proteolytical degradation while not impairing activity. An example is given in the paper "Tritiated D-ala<sup>1</sup>-Peptide T Binding", Smith, C.S. et al, Drug Development Res. 15, pp. 371-379 (1988). Secondly, cyclic structure for stability, such as N to C interchain imides and lactames (Ede et al in Smith and Rivier (Eds) "Peptides: Chemistry and Biology", Escom, Leiden (1991), p268-270), and sometimes also receptor binding may be enhanced by forming cyclic analogues. An

- 12 -

example of this is given in "Conformationally restricted  
thymopentin-like compounds", U.S. pat. 4,457,489 (1985),  
Goldstein, G. et al. Finally, the introduction of  
ketomethylene, methylsulfide or retroinverse bonds to  
5 replace peptide bonds, i.e. the interchange of the CO and  
NH moieties may both greatly enhance stability and  
potency. An example of the latter type is given in the  
paper "Biologically active retroinverse analogues of  
thymopentin", Sisto A. et al in Rivier, J.E. and Marshall,  
10 G.R. (eds.) "Peptides, Chemistry, Structure and Biology",  
Escom, Leiden (1990), p.722-773.

The peptides of the invention can be synthesized by  
various methods which are known in principle, namely by  
chemical coupling methods (cf. Wunsch, E.: "Methoden der  
15 organischen Chemie", Volume 15, Band 1 + 2, Synthese von  
Peptiden, Thieme Verlag, Stuttgart (1974), and Barrany, G.;  
Merrifield, R.B: "The Peptides", eds. E. Gross,  
J. Meienhofer., Volume 2, Chapter 1, pp. 1-284, Academic  
Press (1980)), or by enzymatic coupling methods  
20 (cf. Widmer, F., Johansen, J.T., Carlsberg Res. Commun.,  
Volume 44, pp. 37-46 (1979), and Kullmann, W.: "Enzymatic  
Peptide Synthesis", CRC Press Inc., Boca Raton, Florida  
(1987), and Widmer, F., Johansen, J.T. in "Synthetic  
Peptides in Biology and Medicine:", eds., Alitalo, K.,  
25 Partanen, P., Vattieri, A., pp. 79-86, Elsevier, Amsterdam  
(1985)), or by a combination of chemical and enzymatic  
methods if this is advantageous for the process design and  
economy.

It will be seen that one of the alternatives embraced  
30 in the general formula set out above is for a cysteine  
residue to be positioned at both the amino and carboxy  
terminals of the peptide. This will enable the cyclisation  
of the peptide by the formation of di-sulphide bond.

It is intended that such modifications to the peptide  
35 of the present invention which do not result in a decrease  
in biological activity are within the scope of the present  
invention.

- 13 -

As would be recognized by those skilled in the art there are numerous examples to illustrate the ability of anti-idiotypic (anti-Ids) antibodies to an antigen to function like that antigen in its interaction with animal cells and components of cells. Thus, anti-Ids to a peptide hormone antigen can have hormone-like activity and interact specifically with the receptors to the hormone. Conversely, anti-Ids to a receptor can interact specifically with a mediator in the same way as the receptor does. (For a review of these properties see: Gaulton, G.N. and Greane, M.I. 1986. Idiotypic mimicry of biological receptors, *Ann. Rev. Immunol.* 4, 253-280; Sege, K and Peterson, P.A., 1978. Use of anti-idiotypic antibodies as cell surface receptor probes. *Proc. Natl. Acad. Sci. U.S.A.* 75, 2443-2447).

As might be expected from this functional similarity of anti-Id and antigen, anti-Ids bearing the internal image of an antigen can induce immunity to such an antigen. (This nexus is reviewed in Hiernaux, J.R. 1988. Idiotypic vaccines and infectious diseases. *Infect. Immun.* 56, 1407-1413.)

As will be appreciated by persons skilled in the art from the disclosure of this application it will be possible to produce anti-idiotypic antibodies to the peptide of the present invention which will have similar biological activity. It is intended that such anti-idiotypic antibodies are included within the scope of the present invention.

Accordingly, in a fourth aspect the present invention consists in an anti-idiotypic antibody to the peptide of the first aspect of the present invention, the anti-idiotypic antibody being capable of abrogating TNF and/or LPS toxicity.

The individual specificity of antibodies resides in the structures of the peptide loops making up the Complementary Determining Regions (CDRs) of the variable

- 14 -

domains of the antibodies. Since in general, the amino acid sequences of the CDR peptide loops of an anti-Id are not identical to or even similar to the amino acid sequence of the peptide antigen from which it was originally derived, it follows that peptides whose amino acid sequence is quite dissimilar, in certain contexts can take up a very similar three-dimensional structure. The concept of this type of peptide, termed a "functionally equivalent sequence" or mimotope by Geyson is familiar to those expert in the field. (Geyson, H.M. et al 1987. Strategies for epitope analysis using peptide synthesis. J. Immun. Methods. 102, 259-274).

Moreover, the three-dimensional structure and function of the biologically active peptides can be simulated by other compounds, some not even peptidic in nature, but which mimic the activity of such peptides. This field of science is summarised in a review by Goodman, M. (1990). (Synthesis, spectroscopy and computer simulations in peptide research. Proc. 11th American Peptide Symposium published in Peptides-Chemistry, Structure and Biology pp 3-29. Ed Rivier, J.E. and Marshall, G.R. Publisher ESCOM.)

As will be recognized by those skilled in the art, armed with the disclosure of this application, it will be possible to produce peptide and non-peptide compounds having the same three-dimensional structure as the peptide of the present invention. These "functionally equivalent structures" or "peptide mimics" will react with antibodies raised against the peptide of the present invention and may also be capable of abrogating TNF toxicity. It is intended that such "peptide mimics" are included within the scope of the present invention.

Accordingly, in a fifth aspect the present invention consists in a compound the three-dimensional structure of which is similar as a pharmacophore to the three-dimensional structure of the peptide of the first aspect



- 15 -

of the present invention, the compound being characterized in that it reacts with antibodies raised against the peptide of the first aspect of the present invention and that the compound is capable of abrogating TNF and/or LPS toxicity.

More detail regarding pharmacophores can be found in Bolin et al. p 150, Polinsky et al. p 287, and Smith et al. p 485 in Smith and Rivier (Eds) "Peptides: Chemistry and Biology", Escom, Leiden (1991).

#### 10 Detailed Description of the Invention

In order that the nature of the present invention may be more clearly understood, the preferred forms thereof will now be described with reference to the following example and accompanying Figures and Tables in which:

15 Fig. 1 shows the amino acid sequence of human TNF $\alpha$  ;

Fig. 2: Effect of TNF ( $\square$ ) and TNF+ Peptide 1 ( $\blacklozenge$ ) on blood glucose levels in malaria primed mice-Peptide 1 abrogates TNF induced hypoglycaemia in malaria primed mice.

Fig. 3: Effect of Peptide 1 on TNF-induced tumour regression.

20 Fig. 4: Effect of Peptide 1 ( $\bullet$ ), peptide 308 ( $\blacktriangledown$ ), peptide 309 ( $\blacksquare$ ), peptide 305 ( $\boxtimes$ ) and peptide 302 ( $\circ$ ) on binding of radiolabelled TNF to TNF receptors on WEHI-164 tumour cells - Peptide 1 does not inhibit binding of TNF to tumour cells.

Fig. 5: Plasma reactive nitrogen intermediate levels in TNF+ Peptide 1 treated malaria primed mice - this shows that induction of RNI by TNF is inhibited by treatment with Peptide 1.

30 Fig. 6 shows the effect on blood glucose levels in mice treated with PBS ( $\square$ ); TNF alone ( $\blacklozenge$ ); TNF + Peptide 1 ( $\blacksquare$ ) and TNF + Peptide 2 ( $\circ$ ).

Fig. 7 shows the effect of Peptide 1 on TNF-induced decrease in blood glucose levels in mice administered with 35 200 $\mu$ g TNF.

- 16 -

Fig. 8 shows the effect of Peptide 1 on TNF-induced  
~ decrease in blood glucose levels in ascites tumour-bearing mice.

Fig. 9 shows the effect of Peptide 1 on TNF-induced  
5 weight loss in ascites tumour-bearing mice.

Fig. 10 shows the effect of peptides on LPS toxicity in Meth A ascites tumour-bearing mice (10 animals per group scored positive if 7 or more survive);

Fig. 11 shows the effect of peptides on LPS toxicity  
10 in Meth A ascites tumour-bearing mice (10 animals per group scored positive if 7 or more survive);

Fig. 12 shows the effect of peptides on TNF toxicity in Meth A ascites tumour-bearing mice (each group contains 20 animals: scored positive if 7 or more survived);

15 Fig. 13 shows the effect of peptides on TNF toxicity in Meth A ascites tumour-bearing mice (each group contains 20 animals: scored positive if 10 or more survived);

Fig. 14 shows effect of peptides on TNF toxicity in D-galactosamine sensitized mice (each group contains 10  
20 animals: scored positive if 6 or more survive).

Fig. 15 shows the effect of peptides on direct induction of chemiluminescence by TNF on human neutrophils;

Fig. 16 shows inhibition of TNF priming of human neutrophils by Peptide 21;

25 Fig. 17 shows inhibition of TNF priming of human neutrophils by Peptide 19;

Fig. 18 shows inhibition of LPS stimulation of neutrophils by Peptide 19;

Fig. 19 shows dose-dependent effects of Peptide 9 on  
30 TNF-induced chemiluminescence;

Fig. 20 shows effect of peptide 2 on human TNF priming of human neutrophils;

Fig. 21 shows inhibition of LPS-induced chemiluminescence response of human neutrophils by Peptid  
35 21; and

Fig. 22 shows inhibition of TNF priming of human neutrophils by Peptide 21.

- 17 -

### Production of Peptides

#### Synthesis of Peptides Using the FMOC-Strategy

Peptides (1-6, 9-18, 22-25, 27-29, 35, 36, 39, 40 Table 3) were synthesized on the continuous flow system as provided by the Milligen synthesizer Model 9050 using the standard Fmoc-polyamide method of solid phase peptide synthesis (Atherton et al, 1978, J.Chem. Soc. Chem. Commun., 13, 537-539).

For peptides with free carboxyl at the C-terminus, the solid resin used was PepSyn KA which is a polydimethylacrylamide gel on Kieselguhr support with 4-hydroxymethylphenoxyacetic acid as the functionalised linker (Atherton et al., 1975, J.Am.Chem.Soc 97, 6584-6585). The carboxy terminal amino acid was attached to the solid support by a DCC/DMAP-mediated symmetrical-anhydride esterification.

For peptides with carboxamides at the C-terminus, the solid resin used was Fmoc-PepSyn L Am which is analogous polyamides resin with a Rink linker, p-[(R,S)- $\alpha$ [1-(9H-fluoren-9-yl)-methoxyformamido]-2, 4-dimethoxybenzyl]-phenoxyacetic acid (Bernatowicz et al, 1989, Tet.Lett. 30, 4645). The synthesis starts by removing the Fmoc-group with an initial piperidine wash and incorporation of the first amino acid is carried out by the usual peptide coupling procedure.

The Fmoc strategy was also carried out in the stirred cell system in synthesis of peptides (33,34,37,38) where the Wang resin replaced the Pepsyn KA.

All Fmoc-groups during synthesis were removed by 20% piperidine/DMF and peptide bonds were formed either of the following methods except as indicated in Table 1:

1. Pentafluorophenyl active esters. The starting materials are already in the active ester form.
2. Hydroxybenzotriazol esters. These are formed in situ either using Castro's reagent, BOP/NMM/HOBt (Fournier et al, 1989, Int.J.Peptide Protein Res., 33, 133-139) or

- 18 -

using Knorr's reagent, HBTU/NMM/HOBt (Knorr et al, 1989, Tet.Lett., 30, 1927).

Side chain protection chosen for the amino acids was removed concomitantly during cleavage with the exception of Acn on cysteine which was left on after synthesis. Intramolecular disulphide bridges where needed are then formed by treating the Acn protected peptide with iodine/methanol at high dilution.

TABLE 1

	<u>Amino Acid</u>	<u>Protecting Group</u>	<u>Coupling Method</u>
10	Arg	Pmc	HOBt or OPfp
	Asp	OBu	HOBt or OPfp
	Cys	Acn	HOBt or OPfp
	Glu	OBu	HOBt or OPfp
15	His	Boc or Trt	HOBt or OPfp
	Lys	But	HOBt or OPfp
	Ser	But	HOBt only
	Thr	But	HOBt only
	Tyr	But	HOBt or OPfp
20	Asn	none	OPfp only
	Gln	none	OPfp only

#### Cleavage Conditions

Peptides were cleaved from the PepSyn KA and PepSyn K Am using 5% water and 95% TFA where Arg(Pmc) is not present. Where Arg(Pmc) is present a mixture of 5% thioanisole in TFA is used. The cleavage typically took 3 h at room temperature with stirring. Thioanisole was removed by washing with ether or ethyl acetate and the peptide was extracted into an aqueous fraction. Up to 30% acetonitrile was used in some cases to aid dissolution. Lyophilization of the aqueous/acetonitrile extract gave the crude peptide.

Peptides from the Wang resin were cleaved using 5% phenol, 5% ethanedithiol and 90% TFA for 16 h at ambient temperature with stirring. Thioanisole was removed by

- 19 -

washing with ether or ethyl acetate and the peptide was extracted into an aqueous fraction. Up to 30% acetonitrile was used in some cases to aid dissolution. Lyophilization of the aqueous/acetonitrile extract gave the crude peptide.

Peptides from the Wang resin were cleaved using 5% phenol, 5% ethanedithiol and 90% TFA for 16 h at ambient temperature with stirring.

#### Purification

Crude peptide is purified by reverse phase chromatography using either a C4 or C18 column and the Buffer system: Buffer A - 0.1% aqueous TFA, Buffer B - 80% Acetonitrile and 20% A.

#### N-Terminal Acetylation

The peptide resin obtained after the synthesis (with Fmoc removed in the usual manner was) placed in a 0.3 MDMF solution of 10 equivalents of Ac-OHSu for 60 minutes. The resin was filtered, washed with DMF, CH<sub>2</sub>Cl<sub>2</sub>, ether and used in the next step.

#### Cyclization

The purified and lyophilized bis-S-(acetamidomethyl) cysteine peptide (100-400 mg) was dissolved in 5 mls of methanol containing 1 ml of acetic acid. This was added dropwise to a 1 litre methanol solution containing 1 g of iodine.

After 2 h reaction, the excess iodine was removed by addition of a dilute sodium thiosulfate solution until the colour turns to a pale yellow, methanol was removed in vacuo at room temperature and the concentrated solution was finally completely decolourised with dropwise addition of sodium thiosulfate and applied immediately onto a preparatively reverse phase chromatography column.

#### Synthesis of Peptides using the Boc-Strategy

Syntheses of these peptides were carried out on the ABI 430A instrument using polystyrene based resins. For peptide with C-terminal acids, the appropriate Merrifield

- 20 -

resin Boc-amino acid-O-resin or the 100-200 mesh PAM resin is used (7, 8, 19-21, 26, 31). Peptides with C-terminal amides are synthesized on MBHA resins (32, 33).

Couplings of Boc-amino acids (Table 2) were carried out either using symmetrical anhydride method or a HOBt ester method mediated by DCC or HTBU.

TABLE 2

	<u>Amino Acid</u>	<u>Protecting Group</u>	<u>Coupling Method</u>
	Arg	Tos	HOBt or S.A.
10	Asp	Cxl, OBzl	HOBt or S.A.
	Cys	4-MeBzl	HOBt or S.A.
	Glu	Cxl	HOBt or S.A.
	His	Dnp, Bom	HOBt or S.A.
	Lys	2-Clz	HOBt or S.A.
15	Ser	Bzl	HOBt or S.A.
	Thr	Bzl	HOBt or S.A.
	Tyr	Br-Z	HOBt or S.A.
	Asn	Xan	HOBt or S.A.
	Gln	none	HOBt only

20

Cleavage

Peptides were cleaved in HF with p-cresol or anisole as scavenger for up to 90 min. For His with Dnp protection, the resin required pre-treatment with mercaptoethanol:DIPEA:DMF (2:1:7), for 30 min. After removal of scavengers by ether wash, the crude peptide is extracted into 30% acetonitrile in water.

N-Terminal Acetylation

Acetylation was achieved by treating the deblocked resin with acetic anhydride in DMF solution.

TABLE 3

	<u>No</u>	<u>hTNF</u>	<u>Sequence</u>
	1	1-18	VAL ARG SER SER SER ARG THR PRO SER ASP LYS PRO VAL ALA HIS VAL VAL ALA
35	2	6-18	ARG THR PRO SER ASP LYS PRO VAL ALA HIS VAL VAL ALA

- 21 -

3	2-15	ART SER SER SER ARG THR PRO SER ASP LYS PRO VAL ALA HIS
4	1-26	VAL ARG SER SER SER ARG THR PRO SER ASP LYS PRO VAL ALA HIS VAL VAL ALA ASN PRO GLN ALA GLU GLY GLN LEU
5		
5	10-18	ASP LYS PRO VAL ALA HIS VAL VAL ALA
6	15-22	HIS VAL VAL ALA ASN PRO GLN ALA
7	6-16	ARG THR PRO SER ASP LYS PRO VAL ALA HIS VAL
10	8 6-17	ARG THR PRO SER ASP LYS PRO VAL ALA HIS VAL VAL
	9 8-16	PRO SER ASP LYS PRO VAL ALA HIS VAL
	10 8-15	PRO SER ASP LYS PRO VAL ALA HIS
	11 8-15	PRO SER ASP LYS PRO VAL ALA
15	12 8-13	PRO SER ASP LYS PRO VAL
	13 7-18	THR PRO SER ASP LYS PRO VAL ALA HIS VAL VAL ALA
	14 8-18	PRO SER ASP LYS PRO VAL ALA HIS VAL VAL ALA
20	15 9-18	SER ASP LYS PRO VAL ALA HIS VAL VAL ALA
	16 11-18	LYS PRO VAL ALA HIS VAL VAL ALA
	17 12-18	PRO VAL ALA HIS VAL VAL ALA
	18 12-18	Ac PRO VAL ALA HIS VAL VAL ALA NH2
	19 6-18	ARG THR PRO SER <u>ALA</u> LYS PRO VAL ALA HIS VAL VAL ALA
25		
	Ala(10)	
	20 6-18	ARG THR PRO SER ASP <u>ALA</u> PRO VAL ALA HIS VAL VAL ALA
	Ala(11)	
30	21 6-18	ARG THR PRO SER <u>LYS</u> <u>ASP</u> PRO VAL ALA HIS VAL VAL ALA
	Lys(10)	
	Asp(11)	
	22 1-18	VAL ARG SER SER SER ARG THR PRO SER ASP LYS PRO VAL ALA <u>ARG</u> VAL VAL ALA
35		
	Arg(15)	

- 22 -

	23	1-18	VAL ARG SER SER SER ARG THR PRO SER ASP
		GLN(15)	LYS PRO VAL ALA GLN VAL VAL ALA
	24	1-18	VAL ARG SER SER SER ARG THR PRO SER ASP
		Leu(14)	LYS PRO VAL LEU HIS VAL VAL ALA
5	25	1-18	VAL ARG SER SER SER ARG THR PRO SER ASP
			LYS PRO VAL VAL HIS VAL VAL ALA
		Val(14)	
	26	6-26	ARG THR PRO SER ASP LYS PRO VAL ALA HIS
			VAL VAL ALA ASN PRO GLN ALA GLU GLY GLN
10			LEU
	27	1-16	VAL ARG SER SER SER ARG THR PRO SER ASP
			LYS PRO VAL ALA HIS VAL
	28	1-10	VAL ARG SER SER SER ARG THR PRO SER ASP
	29	8-14	Ac PRO SER ASP LYS PRO VAL ALA NH2
15	30	6-16	Ac ARG THR PRO SER ASP LYS PRO VAL ALA
			HIS VAL NH2
	31	6-16	ARG THR PRO SER ASP LYS PRO VAL VAL HIS
			VAL
		Val(14)	
20	32	6-16	ARG THR PRO SER ASP LYS PRO VAL ALA HIS
			ALA
		ALA(16)	
	33	6-16	ARG THR PRO SER ASP LYS PRO VAL ALA ALA
			VAL
25		ALA(15)	
	34	6-16	ART THR PRO SER ASP LYS PRO VAL ALA LYS
			VAL
		LYS(15)	
	35	6-16	ARG THR PRO SER ASP LYS PRO VAL ALA ASP
30			VAL
		ASP(15)	
	36	6-16	ARG THR PRO SER ASP LYS PRO VAL ALA D-HIS
			VAL
		D-HIS(15)	
35	275	111-120	ALA LYS PRO TRP TYR GLU PRO ILE TYR LEU



- 23 -

302 43-48 LEU ARG ASP ASN GLN LEU VAL VAL PRO SER  
 SLU GLY LEU TYR LEU ILE  
 303 94-109 LEU SER ALA ILE LYS SER PRO LYS GLN ARG  
 GLU THR PRO GLU GLY ALA  
 5 304 63-83 LEU PHE LYS GLY GLN GLY CYS PRO SER THR  
 HIS VAL LEU LEU THR HIS THR ILE SER ARG  
 ILE  
 305 132-150 LEU SER ALA GLU ILE ASN ARG PRO ASP TYR  
 LEU ASP PHE ALA GLU SER GLY GLN VAL  
 10 306 13-26 VAL ALA HIS VAL VAL ALA ASN PRO GLN ALA  
 GLU GLY GLN LEU  
 307 22-40 ALA GLU GLY GLN LEU GLN TRP LEU ASN ARG  
 ARG ALA ASN ALA LEU LEU ALA ASN GLY  
 308 54-68 GLY LEU TYR LEU ILE TYR SER SLN VAL LEU  
 15 PHE LYS GLY GLN GLY  
 309 73-94 HIS VAL LEU LEU THR HIS THR ILE SER ARG  
 ILE ALA VAL SER TYR GLN THR LYS VAL ASN  
 LEU LEU  
 323 79-89 THR ILE SER ARG ILE ALA VAL SER TYR GLN  
 20 THR  
 347 132-157 LEU SER ALA GLU ILE ASN ARG PRO ASP TYR  
 LEU ASP PHE ALA GLU SER GLY GLN VAL TYR  
 PHE GLY ILE ILE ALA LEU

#### Endothelial Cell Clotting Assays

25 Endothelial cell procoagulant activity (PCA)  
 induction by  $\text{TNF}\alpha$  was determined using bovine aortic  
 endothelial cells (BAE) according to the procedure of  
 Bevilacqua et al., 1986 PNAS 83, 4522 with the following  
 modifications: BAE cells were propagated in McCoys 5A  
 30 medium supplemented with 10% FCS, penicillin, streptomycin  
 and L-glutamine in standard tissue culture flasks and  
 24-well dishes.  $\text{TNF}\alpha$  treatment of culture ( $3\mu\text{g/ml}$ ) was  
 for 4 hours at  $37^{\circ}\text{C}$  in the presence of growth medium  
 after which the cells were washed and scrape-harvested  
 35 before being frozen, thawed and sonicated. Total cellular  
 PCA was determined in a standard one-stage clotting assay

- 24 -

using normal donor platelet poor plasma to which 100 $\mu$ l of  $\sim$  CaCl<sub>2</sub> and 100 $\mu$ l of cell lystate was added. Statistical significance was determined by unpaired t-test.

#### Neutrophil Activation Studies

5 In these experiments, neutrophils were prepared from blood of healthy volunteers by the rapid single step method (Kowanko and Ferrante 1987 Immunol 62, 149). To 100 $\mu$ l of  $5 \times 10^6$  neutrophils/ml was added 100 $\mu$ l of either 0, 10, 100 $\mu$ g of peptide/ml and 800 $\mu$ l of  
10 lucigenin (100 $\mu$ g). The tubes were immediately placed into a light proof chamber (with a 37°C water jacket incubator) of a luminometer (model 1250; LKB Instruments, Wallac, Turku, Finland). The resultant light output (in millivolts was recorded). The results are recorded as the  
15 maximal rate of chemiluminescence production.

Effects of peptides on neutrophil chemiluminescence induced by either TNF or LPS: Neutrophils of 96-99% purity and >99% viability were prepared from blood of normal healthy volunteers by centrifugation (400g for 30  
20 min) through Hypaque-Ficoll medium of density 1.114. Following centrifugation the neutrophils formed a single band above the erythrocytes and 1 cm below the mononuclear leukocyte band. These were carefully recovered and washed in medium 199. To assess the lucigenin-dependent  
25 chemiluminescence response 100ul of  $5 \times 10^6$  neutrophils/ml was added 100ul of either 0, 1, 10, 100ug of peptide/ml and TNF or LPS and 800ul of lucigenin (100ug). The tubes were immediately placed into a light proof chamber with a 37°C water jacket incubator of a  
30 luminometer. The resultant light output (in millivolts) was recorded. The results are recorded as the maximal of chemiluminescence production. In experiments which examined the ability of the peptides to prime for the response to fMLP, 100ul of  $5 \times 10^5$  neutrophils /ml  
35 preincubated in peptide and LPS or TNF for 20 mins was

- 25 -

add d to 100ul of diluent or fMLP ( $5 \times 10^{-6}$  M) before the addition of 700ul of lucigenin (100ug). The chemiluminescence was measured as above. Neutrophils from at least three individuals were used in triplicate determinations of anti-TNF or LPS activity. Results were deemed positive if at least 50% inhibition of chemiluminescence was obtained in at least two thirds of cases.

#### WEH1-164 Cytotoxicity

10 Bioassay of recombinant TNF activity was performed according to the method described by Espevik and Nissen-Meyer. (Espevik and Nissen-Meyer 1986 J. Immunol. Methods 95 99-105)

#### Tumour Regression Experiments

15 Subcutaneous tumours were induced by the injection of approximately  $5 \times 10^5$  WEH1-164 cells. This produced tumours of diameters of 10 to 15mm approximately 14 days later. Mice were injected i.p. with recombinant human TNF (10µg and 20µg) and peptide (1mg) for four consecutive  
20 days. Control groups received injections of PBS. Tumour size was measured daily throughout the course of the experiment. Statistical significance of the results was determined by unpaired Student T-test.

#### Radioreceptor assays

25 WEH1-164 cells grown to confluency were scrape harvested and washed once with 1% bovine serum albumin in Hanks balanced salt solution (HBSS, Gibco) and used at  $2 \times 10^6$  cells pre assay sample. For the radioreceptor assay, the cells were incubated with varying amounts of  
30 either unlabelled TNFα(1- $10^4$  ng per assay sample) or peptide (0- $10^5$  ng per assay sample) and  $^{125}$ I-TNF (50,000cpm) for 3 hours at 37°C in a shaking water bath. At the completion of the incubation 1ml of HBSS/BSA was added to th WEH1-164 cells, the cells spun and the  
35 bound  $^{125}$ I in the cell pellet counted. Specific binding

- 26 -

was calculated from total binding minus non-specific binding of triplicate assay tubes. 100% specific binding corresponded to 1500 cpm.

#### In Vivo Studies of TNF Toxicity

5 Mice were administered with either TNF (200µg), Peptide 1 (10mg) and TNF (200µg)+Peptide 1 (10mg) via intravenous injection. Blood glucose levels and appearance of the animals was evaluated at 15, 30, 60, 120, 180 minutes after injection. Appearance parameters  
10 which were evaluated included ruffling of fur, touch sensitivity, presence of eye exudate, light sensitivity and diarrhoea.

#### Infection of mice with malaria parasites and treatment with TNF+ Peptide 1

15 All the mice used were male, CBA/CaH strain and 6-8 weeks old. *P. vinkei vinkei* (Strain V52, from F.E.G. Cox, London) has undergone several serial passages in CBA mice, after storage in liquid nitrogen, before use in these experiments. Infections were initiated by intraperitoneal  
20 injection of  $10^6$  parasitized erythrocytes. Mice were treated with TNF(7µg) ± peptide (8.3 mg) administered iv.

#### Assays for blood glucose

Nonfasting blood glucose levels were determined on a Beckman Glucose Analyzer 2 (Beckman Instruments) or on a  
25 Exectech blood glucose sensor (Clifford Hallam Pty. Ltd).

#### Reactive Nitrogen Intermediates (RNI)

RNI levels in blood were determined by the method of Rockett et al (1991) in-vivo induction of TNF, LT and IL-1 implies a role for nitric oxide in cytokine-induced  
30 malarial cell-mediated immunity and pathology. J. Immunol. in press.

TNF and LPS Lethality Experiments: balb/C or balbC x swiss F1 mice carrying Meth A ascites tumours elicited by prior I.P. inoculation of 0.5µl pristane 7 days before  
35 I.P. injection of tumour cells. Nine to ten days after

- 27 -

inoculation with the tumour cells 25 ug of human recombinant TNF was subcutaneously administered and a short time later 1mg of either test peptide, bovine serum albumen, phosphate buffered saline or neutralizing anti-TNF MAb 47 was administered at a separate subcutaneous site. The number of surviving animals was then observed at 18 hours and 24 hours post TNF treatment. In experiments which assessed the effects of 1-related peptides on on LPS lethality the mice were administered 500ug E.coli LPS and peptide or other treatment in a similar manner. In LPS experiments polymyxin B, an LPS inhibitor, replaced MAb 47 as a positive control. The number of animals surviving was assessed at intervals up to 64 hours after LPS challenge.

15 Experiments in D-galactosamine sensitized mice: Female Bablb/C mice were co-injected intraperitoneally with 16 mg D-galactosamine and 2ug human recombinant TNF. The mice were then injected subcutaneously with either test peptide, phosphate buffered saline or neutralizing anti-TNF monoclonal antibody 47. The number of surviving animals was assessed at intervals up to 48 hours after TNF challenge.

#### RESULTS

The results obtained with each of the peptides are summarised in Table 4. A single \* indicates heightened activity in that test whilst a double \*\* indicates activity at low concentrations of peptide but not high concentrations.

- 28 -

TABLE 4

PEPTIDE	IN VIVO			IN VITRO NEUTROPHIL			
	TNF TOXICITY		LPS TOXICITY	DIRECT	TNF	LPS	
	METH A	D-GAL	METH A		PRIMING	DIRECT	PRIMING
1	+	+	+	+	+	+	+
2	++	+	+	++			
8	-		-	+			
9	-		-	+++			
10	++	-	-	+			
11	-			-			
12	+			-			
16	-			-			
17	-		+	-			
13	-		-	+			
14	-		+	+			
15	-		-	-			
18	-	-					
19	+		+	+	+	+	+
20	-		-	-			
21	++		+	+	+	+	+
22		+	+	+	+		
23	+	+	+	+			
24	-		-	-			
25	+/-		-	+			
26	-		-	+			
4	-			+			
5	-		-	+			
6	-			-			
3	-						
28	+	-	+				
29	-	-	+				
30	++	+	+				
31	+	+	-				
32	-		-				
33	-		++				
34	-		++				
36	-		-				
35	+		+				
27	-		-				
7	-		+	++			

- 29 -

TNF administered at a dose of 200µg was found to be toxic in mice according to the parameters studied. In particular, blood glucose levels had fallen by 120 minutes (Fig 7) Peptide 1 alone in 2 of the 3 mice studied did not reduce blood glucose levels. Mouse 1 in this group recovered normal blood glucose levels within by 180 minutes. Mice in the group treated with a combination of TNF and Peptide 1 showed no reduction in blood glucose levels at 120 min and a small decrease at 180 min.

10 As shown in Fig. 6, 10µg of Peptide 2 given to mice treated with 200µg of recombinant human TNF abrogated TNF toxicity as indicated by the inhibition of blood glucose changes evident in mice treated with TNF alone.

When general appearance of treated mice was considered it was noted that all 3 TNF only treated mice had ruffled fur, touch sensitivity and light sensitivity. One mouse in this group also had diarrhoea. Mice treated with Peptide 1 alone showed only slight touch sensitivity with one mouse showing slight ruffling of the fur at 180 mins. Mice treated with a combination of TNF and Peptide 1 showed ruffling of the fur and slight touch sensitivity at 180 mins but failed to show either light sensitivity or onset of diarrhoea. In addition, Peptide 1 and related peptides prevented death in acute models of TNF lethality (Figs. 12 & 13).

Peptide 1 failed to either activate the respiratory burst of human neutrophils (Table 5) or to induce procoagulant activity on bovine aortic endothelial cells, and hence is free of these negative aspects of TNF activity in acute or chronic inflammation. However, Peptide 1 and related peptides inhibited both the TNF and LPS-induced respiratory burst of human neutrophils (Figs. 15, 19, 18, 21). Further, several peptides inhibited priming of the neutrophil response to a bacterially-derived peptid EMLP (Figs. 16, 17, 20, 22).

- 30 -

TABLE 5

Peptide	Concentration ug/10 <sup>6</sup> cells)				
	0	1	10	100	500
5					
275	1.02	0.99	0.69	0.43	0.80
1	0.34	0.93	0.74	0.55	1.10
302	0.37	0.15	0.18	0.29	
303	0.37	0.22	0.17	0.22	
10 304	0.37	0.18	0.43	2.56	2.76
305	0.37	0.27	0.36	0.24	
306	0.37	0.27	0.35	0.23	
307	0.37	0.35	0.37	0.42	
323	0.37	0.23	0.17	0.47	
15 308	0.37	0.91	1.80	49.52	
309	0.37	0.38	0.98	13.44	

Results are expressed as mV of lucigenin dependent  
 20 chemiluminescence and represent peak of response i.e. the  
 maximal cell activity attained.

The results shown in Fig. 3 clearly show one of the  
 desirable effects of TNF $\alpha$ , i.e. tumour regression, is  
 unaffected by Peptide 1. Further, Peptide 1 does not  
 25 inhibit binding of TNF to tumour cell receptors (Fig 4).  
 Table 6 indicates that Peptide 1 is devoid of intrinsic  
 anti-tumour activity. The ability of Peptide 1 to prevent  
 high plasma RNI levels in TNF $\alpha$  treated malaria primed  
 mice is also strongly indicative of the therapeutic  
 30 usefulness of this peptide (Fig 5). Peptide 1 also  
 inhibits the TNF-induced decrease in blood glucose levels  
 evident in mice treated with TNF alone (Fig 2). Further  
 in the experiments involving mice infected with malaria  
 parasites; of the three mice treated with TNF $\alpha$  alone one  
 35 died and the other two were moribund. In contrast in th



- 31 -

group of three mice treated with TNF $\alpha$  and Peptid 1 all survived and non were moribund. This very marked result also strongly indicates the potential usefulness of this peptide as a therapeutic.

5 Peptide 1 inhibits not only the TNF-induced hypoglycaemia in sensitized mice but also in ascites tumour-bearing mice (Fig 8). Further, tumour-bearing mice treated with TNF + Peptide 1 fail to develop the cachexia or weight loss associated with TNF treatment (Fig 9).

10 As will be seen from the above information the peptide of the present invention are capable of abrogating TNF and/or LPS toxicity in vivo and neutrophil activation by LPS or TNF in vitro. This peptide has utility in the treatment of numerous disease states which are due to the deleterious effects of TNF and/or LPS.

TABLE 6

In vitro cytotoxicity of TNF and synthetic TNF peptides on WEHI 164 fibrosarcoma cells

	<u>TNF/PEPTIDE</u>	<u>% VIABLE CELLS*</u>
20	TNF#	26.6
	275+	100
	1	100
	302	48.7
	304	100
25	305	72.7
	306	100
	307	100
	308	42.2
	309	92.8

30

---

\* %Viability was determined by comparison with untreated control cells. Results shown are the means of quadruplicate determinations.

35 # TNF was at 50 units per cultur which is equivalent to 3ug (12ug/ml)

+ Each peptide was tested at 50ug/culture (200ug/ml)


- 32 -

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the  
5 invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

- 33 -

## CLAIMS:-

1. A linear or cyclic peptid of the general formula:-



in which

- 5  $X_1$  is null, Cys or  $R_1$   
 $X_2$  is null, Cys,  $R_1$  or  $A_1-A_2-A_3-A_4-A_5$   
in which  $A_1$  is Val or Ile or Leu or Met or His  
 $A_2$  is Arg or Cys or His  
 $A_3$  is Ser or Thr or Ala  
10  $A_4$  is Ser or Thr or Ala  
 $A_5$  is Ser or Thr or Ala  
 $X_3$  is Cys,  $R_1$  or  $A_6-A_7$   
in which  $A_6$  is Arg or Cys or His or Absent  
 $A_7$  is Thr or Ser or Ala  
15  $X_4$  is Cys,  $R_1$  or  $A_8-A_9$   
in which  $A_8$  is Pro or an  $N\alpha$ -alkylamino acid  
 $A_9$  is Ser or Thr or Ala  
 $X_5$  is Cys,  $R_1$  or  $A_{10}$   
in which  $A_{10}$  is Asp or Ala or Cys or Glu or Gly  
20 or Arg or His  
 $X_6$  is Cys,  $R_2$  or  $A_{11}-A_{12}-A_{13}$   
in which  $A_{11}$  is absent or Cys or Arg or His or  
Asp or Glu  
 $A_{12}$  is Pro or an  $N\alpha$ -alkylamino acid  
25  $A_{13}$  is Val or Ile or Phe or Tyr or Trp  
or His or Leu or His or Met  
 $X_7$  is null, Cys,  $R_2$  or  $A_{14}-A_{15}$   
in which  $A_{14}$  is Ala or Val or Gly or Ile or Phe  
or Trp or Tyr or Leu or His or Met  
30  $A_{15}$  is absent or His or Arg or Glu or  
Asn or Ala or Lys or Asp or Phe or Tyr or  
Tap or Glu or Gln or Ser or Thr or Gly  
 $X_8$  is null, Cys,  $R_2$ ,  $A_{16}$ ,  $A_{16}-A_{17}$ ,  $A_{16}-A_{17}-A_{18}$  or  
 $A_{16}-A_{17}-A_{18}-A_{19}-A_{20}-A_{21}-A_{22}-A_{23}-A_{24}-A_{25}-A_{26}$

- 34 -

in which      $A_{16}$  is Val or Ile or Leu or Met or His  
               $A_{17}$  is Val or Ile or Leu or Met or His  
               $A_{18}$  is Ala or Gly  
               $A_{19}$  is Asp or Glu  
5               $A_{20}$  is Pro or an  $\alpha$ -alkylamino acid  
               $A_{21}$  is Gln or Asn  
               $A_{22}$  is Ala or Gly  
               $A_{23}$  is Glu or Asp  
               $A_{24}$  is Gly or Aln  
10              $A_{25}$  is Gln or Asn  
               $A_{26}$  is Leu or Ile or Val or Met or His  
 $X_9$  is null, Cys or  $R_2$   
 $R_1$  is R-CO, where R is H, straight, branched or  
cyclic alkyl up to C20, optionally containing double  
15             bonds and/or substituted with halogen, nitro, amino,  
hydroxy, sulfo, phospho or carboxyl groups (which may  
be substituted themselves), or aralkyl or aryl  
optionally substituted as listed for the alkyl and  
further including alkyl, or  $R_1$  is glycosyl,  
20             nucleosyl, lipoyl or  $R_1$  is an L- or D- $\alpha$  amino acid  
or oligomers thereof consisting of up to 5 residues  
 $R_1$  is absent when the amino acid adjacent is an  
unsubstituted desamino-derivative.  
 $R_2$  is  
25              $-NR_{12}R_{13}$ , wherein  $R_{12}$  and  $R_{13}$  are  
independently H, straight, branched or cyclic alkyl,  
aralkyl or aryl optionally substituted as defined for  
 $R_1$  or N-glycosyl or N-lipoyl  
               $-OR_{14}$ , where  $R_{14}$  is H, straight, branched or  
30             cyclic alkyl, aralkyl or aryl, optionally substituted  
as defined for  $R_1$   
              -O-glycosyl, -O-lipoyl or  
              - an L- or D- $\alpha$ -amino acid or a oligamu thereof  
consisting of up to 5 residues  
35             or  $R_2$  is absent, when the adjacent amino acid is a  
decarboxy derivative of cysteine or a homologue  
thereof or the peptide in a N-C cyclic form.

- 35 -

with the proviso that:

- when  $X_6$  is Cys or  $R_2$  then  $X_5$  is  $A_{10}$ ,  $X_4$  is  $A_8-A_9$ ,  
 $X_3$  is  $A_6-A_7$  and  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$   
 when  $X_5$  is Cys or  $R_1$  then  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  
 5  $A_{14}-A_{15}$ ,  $X_8$  is  $A_{16}-A_{17}-A_{18}$  and  $A_{11}$  is absent  
 when  $X_4$  is Cys or  $R_1$  then  $X_5$  is  $A_{10}$ ,  $X_6$  is  
 $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  $A_{14}-A_{15}$  and  $X_8$  is  
 $A_{16}-A_{17}-A_{18}$   
 when  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$  then  $X_8$  is not  $A_{16}$   
 10 when  $X_1$  is null,  $X_2$  is Cys or  $R_1$ ,  $X_3$  is  $A_6-A_7$ ,  $X_4$  is  
 $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  
 $A_{14}-A_{15}$  and  $X_8$  is  $A_{16}$  then  $A_{16}$  is not D-His.  
 $X_1$  is always and only null when  $X_2$  is  $R_1$ , Lys or Null  
 $X_2$  is always and only null when  $X_3$  is Cys or  $R_1$   
 15  $X_3$  is always and only null when  $X_6$  is Cys or  $R_2$   
 $X_7$  is always and only null when  $X_7$  is Cys,  $R_2$  or Null  
 $X_8$  is always and only null when  $X_8$  is Cys,  $R_2$  or Null  
 $X_9$  is always and only null when  $X_8$  is Cys,  $R_2$  or Null  
 when  $X_1$  and  $R_2$  are null,  $X_3$  is  $R_1$ ,  $X_4$  is  
 20  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  $A_{11}-A_{12}-A_{13}$ ,  $X_7$   
 is  $A_{14}-A_{15}$ ,  $X_8$  is  $R_2$  and  $A_{14}$  is Ala and  $A_{15}$  is  
 absent then  $R_1$  is acetyl and  $R_2$  is  $NH_2$ .  
 2. A linear or cyclic peptide as claimed in claim 1 in  
 which:-  
 25  $X_1$  is H,  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$ ,  $X_3$  is  
 $A_6-A_7$ ,  $X_4$  is  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  
 $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  $A_{14}-A_{15}$ ,  $X_8$  is  
 $A_{16}-A_{17}-A_{18}$  and  $X_9$  is OH.  
 3. A linear or cyclic peptide as claimed in claim 1 in  
 30 which:-  
 $X_1$  is null,  $X_2$  is H or Ac,  $X_3$  is  $A_6-A_7$ ,  
 $X_4$  is  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is  
 $A_{11}-A_{12}-A_{13}$ ,  $X_7$  is  $A_{14}-A_{15}$ ,  $X_8$  is  
 $A_{16}-A_{17}-A_{18}$  and  $X_9$  is OH or  $NH_2$ .

- 36 -

4. A linear or cyclic peptide as claimed in claim 1 in which:-

$X_1$  is H,  $X_2$  is  $A_1-A_2-A_3-A_4-A_5$ ,  $X_3$  is  $A_6-A_7$ ,  $X_4$  is  $A_8-A_9$ ,  $X_5$  is  $A_{10}$ ,  $X_6$  is OH and  $X_6$ ,  $X_7$  and  $X_8$  are null.

5. A linear or cyclic peptide as claimed in claim 1 in which the peptide is selected from the group consisting of:-

Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Ala-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Lys-Asp-Pro-Val-Ala-His-Val-Val-Ala;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Arg-Val-Val-Ala;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Gln-Val-Val-Ala;  
 Ac-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-NH<sub>2</sub>;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Ala-Val;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Lys-Val;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val;  
 Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val;  
 Pro-Ser-Asp-Lys-Pro-Val-Ala-His;  
 Pro-Ser-Asp-Lys-Pro-Val;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Val-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala-Asn-Pro-Gln-Ala-Glu-Gly-Gln-Leu;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp;  
 Ac-Pro-Ser-Asp-Lys-Pro-Val-Ala-NH<sub>2</sub>;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Asp-Val;  
 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala-Asn-Pro-Gln-Ala-Glu-Gly-Gln-Leu;  
 Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
 Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val;

- 37 -

Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
Pro-Sir-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;  
Pro-Val-Ala-His-Val-Val-Ala; and  
Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Val-His-Val.

- 5 6. A peptide as claimed in claim 5 in which the peptide is

Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;

Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-His-Val-Val-Ala;

- 10 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp;

Arg-Thr-Pro-Ser-Ala-Lys-Pro-Val-Ala-His-Fal-Val-Ala;

Arg-Thr-Pro-Ser-Lys-Asp-Pro-Val-Ala-His-Val-Val-Ala;

Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Arg-Val-Val-Ala;

- 15 Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Gln-Val-Val-Ala; or

Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala-Asp-Val.

7. A pharmaceutical composition for use in treating subjects suffering from acute or chronic inflammation, the composition comprising a therapeutically effective amount of a peptide as claimed in any one of claims 1 to 6 and a pharmaceutically acceptable sterile carrier.

8. A composition as claimed in claim 7 in which the composition is for administration topically, as a nasal spray, ocularly, intravenously, intraperitoneally, intramuscularly, subcutaneously or for oral delivery.

9. A composition as claimed in claims 7 or 8 in which the composition provides slow release of the active peptide.

- 30 10. A method of treating a subject suffering from acute or chronic inflammation, the method comprising administering to the subject the composition as claimed in any one of claims 7 to 9.

11. A method as claimed in claim 10 in which the subject is suffering from toxic shock, adult respiratory distress

- 38 -

syndrome, hypersensitivity pneumonitis, systemic lupus erythromatosis, cystic fibrosis, asthma, bronchitis, drug withdrawal, schistosomiasis, sepsis, rheumatoid arthritis, acquired immuno-deficiency syndrome, multiple sclerosis, leprosy, malaria, systemic vasculitis, bacterial meningitis, cachexia, dermatitis, psoriasis, diabetes, neuropathy associated with infection or autoimmune disease, ischemia/reperfusion injury, encephalitis, Guillame Barre Syndrome, atherosclerosis, chronic fatigue syndrome, TB, other viral and parasitic diseases and OKT3 therapy.

12. A method of ameliorating or reducing the adverse side effects in a subject receiving cytotoxic drugs, cytokines, immunopotentiating agents, radiation therapy and/or chemotherapy comprising administering to the subject the composition as claimed in any one of claims 7 to 9.

13. An anti-idiotypic antibody to the peptide as claimed in any one of claims 1 to 6, the anti-idiotypic antibody being characterised in that it is capable of abrogating TNF and/or LPS toxicity.

14. A compound the three dimensional structure of which is similar as a pharmacophore to the three dimensional structure of the peptide as claimed in any one of claims 1 to 6, the compound being characterised in that it binds to one or more antibodies raised against the peptides as claimed in any one of claims 1 to 6 and that the compound is capable of abrogating TNF and/or LPS toxicity.



1/22

FIG. 1

**VRSSSRTPSD<sub>10</sub>KPVAHVVA<sub>NP20</sub>QAEGQLQWLN<sub>30</sub>RRA**

**NALLANG<sub>40</sub>VELRDNQLV<sub>50</sub>PSEGLYLIYS<sub>60</sub>QVLFKGQGCP<sub>70</sub>STHVLL**

**THTI<sub>80</sub>SRIAVSYQTK<sub>90</sub>VNLLSAIKSP<sub>100</sub>CQRETREGAE<sub>110</sub>AKPWYEPI**

**YL<sub>120</sub>GGVFQLEKGD<sub>130</sub>RLSAEINRPD<sub>140</sub>YLDFAESGQV<sub>150</sub>YFGIIAL<sub>157</sub>**

2/22




Fig 2

3/22





Fig 3

4/22

FIG. 4



5/22




Fig 5

6/22




Fig 6

7/22




Fig 7

8/22




Fig 8



9/22





Fig 9


10/22

Fig 10




11/22

Fig 11




12/22

Fig 12




13/22

Fig 13



14/22

FIG 14



15/22

FIG 15





FIG 16 16/22





17/22




Fig 17

18/22





FIG 18

19/22

FIG. 19



20/22





FIG 20

21/22

FIG 21



22/22




FIG. 22

## INTERNATIONAL SEARCH REPORT

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)<sup>6</sup>

According to International Patent classification (IPC) or to both National Classification and IPC  
Int. Cl.<sup>8</sup> C07K 7/06, 7/08, 7/10, A61K 37/02

## II. FIELDS SEARCHED

Minimum Documentation Searched<sup>7</sup>

Classification System

Classification Symbols

IPC  
Chem. Abs. online  
CAS online registry

C07K 7/06, 7/08, 7/10, C07C 103/52.  
Keywords: Tumo(u)r Necrosis Factor OR TNF  
PROTEIN SEQUENCE SEARCH

Documentation Searched other than Minimum Documentation,  
to the extent that such Documents are included in the Fields Searched<sup>8</sup>

AU: IPC As Above

III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup>

Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate of the relevant passages <sup>12</sup>	Relevant to Claim No <sup>13</sup>
A	Derwent Abstract Accession no. 90-143138/19, Classes B04 and D16, JP,A, 02-088598 (SOMA G) 28 March 1990 (28.03.90)	1-14
A	Derwent Abstract Accession no. 91-152432/21, Classes B04 and D16, JP,A, 03-087196 (TEIJIN K K) 11 April 1991 (11.04.91)	1-14
A	Derwent Abstract Accession no. 91-145993/20, Classes B04 and D16, JP,A, 03-083587 (TEIJIN K K) 9 April 1991 (09.04.91)	1-14
A	Derwent Abstract Accession no. 91-145992/20, Classes B04 and D16, JP,A, 03-083586 (TEIJIN K K) 9 April 1991 (09.04.91)	1-14

\* Special categories of cited documents : <sup>10</sup>

- "A" Document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

## IV. CERTIFICATION

Date of the Actual Completion of the International Search  
28 August 1992 (28.08.92)

Date of Mailing of this International Search Report  
- 7 SEP 1992 (07.09.92)

International Searching Authority

AUSTRALIAN PATENT OFFICE

Signature of Authorized Officer

A BESTOW



## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE <sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim numbers ..., because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim numbers ..., because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim numbers ..., because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4a

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING <sup>2</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.